



1743 Creekside Drive, Suite 200  
Folsom, CA 95630

530.626.7000 | [www.cgt.global](http://www.cgt.global)

email: [info@cgt.global](mailto:info@cgt.global)

## Fresh Product

### Whole Peripheral Blood Bag Collection, CP2D

Catalog#	PBCPD250F	250 mL
	PBCPD500F	500 mL

## PRODUCT DESCRIPTION

Human Whole Peripheral Blood is collected in a sterile system.

Fresh products have a high viability without the detrimental effects of freezing, thawing, and exposure to cryoprotectants.

Cells were obtained using Institutional Review Board (IRB) approved consent forms and protocols.

## SAMPLE COLLECTION AND PROCESSING

All samples are collected on-site at our Stem Cell Collection Center.

Infectious disease testing for HIV, HBV, and HCV is performed on a sample of donor blood. Only samples with negative results within 90 days of collection are shipped unless approved by the customer. All testing is performed by a CLIA-certified lab.

## FORMAT

Fresh whole peripheral blood is shipped at ambient temperature.

Specific containers and media can also be prepared as requested by the customer.

## STORAGE

Fresh products should be used or processed immediately upon receipt. The warranty only covers items whose specifications are tested at the time they are received.

## CELL COUNTING INSTRUCTIONS

Important: This cell viability/counting step is required to ensure the quantity of cells provided. Be sure to count the cells before washing. Be aware that cell loss is expected and may be up to 30% during wash steps. Recovery rates vary depending on technique.

### Materials

- Cleaned hemocytometer
- Trypan Blue

### Protocol

1. If removing the cell suspension from the vial in which it was shipped, be sure to rinse the vial to collect all of the cells.
2. Gently mix the cell suspension and measure the volume.
3. Make a 1-in-2 dilution with 20  $\mu$ L each of well-mixed cell suspension and Trypan Blue.

4. Load one side of the hemocytometer, being careful not to over- or under-fill the chamber.
5. Count viable (clear, round, bright) and non-viable (blue, irregular shape, dull) cells in the four corner squares. Adjust your dilution if there are more than 100 cells/square.
6. Determine the number of total viable cells in the original sample. One square is equal to 100 nL.

Viability = live cells/all cells

Cell Concentration = Mean cells/square  $\times$  Dilution Factor  $\times$  104

Total Cell Count = Cell Concentration  $\times$  Starting Volume

Total Viable Cell Count = Total Cell Count  $\times$  Viability

## WARNING

This product contains human tissue or other biological material and MUST be handled at Biosafety Level 2 or higher. All biological products should be treated as potentially infectious or contaminated material, even if infectious disease screening reports are negative. Follow universal precautions and wear appropriate personal protective equipment.

## PRODUCT WARRANTY

For our product warranty, please review our Terms and Conditions at [cgt.global/terms-and-conditions/](http://cgt.global/terms-and-conditions/).

FOR RESEARCH USE ONLY. NOT APPROVED FOR HUMAN OR VETERINARY USE IN VIVO.